arXiv:1106.0393v1 [math.RA] 2 Jun 2011

A kind of infinite-dimensional Novikov algebras and its realization *

Liangyun Chen, Yao Ma, Haijun Yu
Department of Mathematics, Northeast Normal University,
Changchun 130024, CHINA

Abstract

In this paper, we construct a kind of infinite-dimensional Novikov algebras and give its realization by hyperbolic sine functions and hyperbolic cosine functions.

Key words: Novikov algebras, left symmetric algebras, adjoining Lie algebras.

AMS Subject Classification: 17A30.

§1 Introduction

The Hamilton operator is an important operator of the calculus of variations. When I. M. Gel'fand and I. Ya. Dorfman [5-7] studied the following operator:

$$H_{ij} = \sum_k c_{ijk} u_k^{(1)} + d_{ijk} u_k^{(0)} \frac{\mathrm{d}}{\mathrm{d}x}, \quad c_{ijk} \in \mathbf{C}, \quad d_{ijk} = c_{ijk} + c_{jik},$$

they gave the definition of Novikov algebras. Concretely, let c_{ijk} be the structural coefficients, a product of $L = L(e_0, e_1, \cdots)$ be $\circ$ such that

$$e_i \circ e_j = \sum c_{ijk} e_k.$$

Then the product is Hamilton operator if and only if $\circ$ satisfies

$$(a \circ b) \circ c = (a \circ c) \circ b$$

---

*Supported by NNSF of China (No.10871057), Natural Science Foundation of Jilin province (No.201115006), Scientific Research Foundation for Returned Scholars Ministry of Education of China and the Fundamental Research Funds for the Central Universities. Corresponding author(L. Chen): chenly640@nenu.edu.cn

$$(a \circ b) \circ c + c \circ (a \circ b) = (c \circ b) \circ a + a \circ (c \circ b).$$

In 1987, E. I. Zel'manov[15] began to study Novikov algebras and proved that the dimension of finite simple Novikov algebras over a field of characteristic zero is one. In algebras, what are paid attention to by mathematician are classifications and structures, but so far we haven't got the systematic theory for general Novikov algebras. In 1992, J. M. Osborn [9-10] had finished the classification of infinite simple Novikov algebras with nilpotent elements over a field of characteristic zero and finite simple Novikov algebras with nilpotent elements over a field of characteristic $p > 0$. In 1995, X. P. Xu [11-14] developed his theory and got the classification of simple Novikov algebras over an algebraically closed field of characteristic zero. C. M. Bai and D. J. Meng [1-3] has serial work on low dimensional Novikov algebras, such as the structure and classification. We construct two kinds of Novikov algebras [4]. Recently, people obtain some properties in Novikov superalgebras [8, 16].

In this paper, we construct a new infinite-dimensional Novikov algebras and give its realization by hyperbolic sine functions and hyperbolic cosine functions.

Definition 1.1 *Let* $(\mathcal{A}, \circ)$ *be an algebra over* $\mathbf{F}$ *such that:*

$$(a, b, c) = (b, a, c), \tag{1.1}$$

$$(a \circ b) \circ c = (a \circ c) \circ b, \quad \forall a, b, c \in \mathcal{A}, \tag{1.2}$$

then $\mathcal{A}$ *is called a Novikov algebra over* $\mathbf{F}$.

Remark 1.2 (1) *Condition* (1.1) *is usually written by*

$$a \circ (b \circ c) - (a \circ b) \circ c = b \circ (a \circ c) - (b \circ a) \circ c. \tag{1.3}$$

(2) *An algebra* $\mathcal{A}$ *is called a left symmetric algebra if it only satisfies* (1.1). *It is clear that left symmetric algebras contain Novikov algebras.*

Remark 1.3 (1) *If* $(\mathcal{A}, \circ)$ *is a left symmetric algebra satisfying*

$$[a, b] = a \circ b - b \circ a, \quad \forall a, b \in \mathcal{A}, \tag{1.4}$$

then $(\mathcal{A}, [,])$ *is a Lie algebra. Usually, it is called an adjoining Lie algebra.*

(2) *Let* $(\mathcal{A}, \cdot)$ *be a commutative algebra, then* $(\mathcal{A}, d_0, \circ)$ *is a Novikov algebra if* d_0 *is a derivation of* $\mathcal{A}$ *with a bilinear operator* $\circ$ *such that*

$$a \circ b = a \cdot d_0(b), \quad \forall a, b \in \mathcal{A}. \tag{1.5}$$

§2 Main results

Lemma 2.1 *Let* $\{ b_0, a_1, b_1, a_2, b_2, \cdots a_n, b_n, \cdots \}$ *be a basis of the linear space* $\mathcal{A}$ *over a field* $\mathbf{F}$ *of characteristic* $p \neq 2$ *satisfying*

$$\begin{cases} a_m a_n = \frac{1}{2}(b_{m+n} - b_{m-n}), \\ b_m b_n = \frac{1}{2}(b_{m+n} + b_{m-n}), \\ a_m b_n = b_n a_m = \frac{1}{2}(a_{m+n} + a_{m-n}), \end{cases} \tag{2.1}$$

where $b_{-m} = b_m$, $a_{-m} = -a_m$. *Then* $\mathcal{A}$ *is a commutative and associative algebra.*

Proof. It is clear that $\mathcal{A}$ is a commutative algebra over $\mathbf{F}$.

$$\begin{aligned} & (a_k, a_n, a_m) = a_k(a_n a_m) - (a_k a_n)a_m \\ = & \frac{1}{2}a_k(b_{m+n} - b_{n-m}) - \frac{1}{2}(b_{k+n} - b_{k-n})a_m \\ = & \frac{1}{4}(a_{k+m+n} + a_{k-m-n} - a_{k+n-m} - a_{k-n+m} \\ & -a_{m+k+n} - a_{m-k-n} + a_{m+k-n} + a_{m-k+n}) \\ = & 0. \end{aligned}$$

Similarly, we have that $(b_k, b_n, b_m) = (a_k, a_n, b_m) = (a_k, b_n, a_m) = (b_k, a_n, a_m) = (b_k, b_n, a_m) = (b_k, a_n, b_m) = (a_k, b_n, b_m) = 0$. Then $(a, b, c) = 0, \forall a, b, c \in \mathcal{A}$. The result follows. □

Corollary 2.2 b_0 *of Lemma 2.1 is a unity of* $\mathcal{A}$.

Lemma 2.3 *Let* $\mathcal{A}$ *be a commutative and associative algebra satisfying Lemma 2.1. Then the following statements hold:*

1) *If* D_0 *is a linear transformation of* $\mathcal{A}$ *such that*

$$\begin{cases} D_0(a_n) = nb_n, & n = 1, 2, 3, \cdots, \\ D_0(b_n) = na_n, & n = 0, 1, 2, \cdots, \end{cases} \tag{2.2}$$

then D_0 *is a derivation of* $\mathcal{A}$.

2) *If* aD_0 *is a linear transformation of* $\mathcal{A}$ *such that*

$$(aD_0)(b) = aD_0(b), \forall a, b \in \mathcal{A}, \tag{2.3}$$

then aD_0 *is a derivation of* $\mathcal{A}$.

3) $\mathcal{D}_1 = \{aD_0 | a \in \mathcal{A}\}$ *is a subalgebra of Lie algebra* Der$\mathcal{A}$.

Proof. (1) We have

$$D_0(a_n a_m) = D_0\left(\frac{1}{2}(b_{n+m} - b_{n-m})\right) = \frac{1}{2}((m+n)a_{n+m} - (n-m)a_{n-m})$$

and

$$\begin{aligned} D_0(a_n)a_m + a_n D_0(a_m) &= nb_n a_m + ma_n b_m \\ &= \frac{n}{2}(a_{n+m} - a_{n-m}) + \frac{m}{2}(a_{n+m} - a_{m-n}) \\ &= \frac{1}{2}((m+n)a_{m+n} - (n-m)a_{n-m}). \end{aligned}$$

So D_0 is a derivation of $\mathcal{A}$.

2) For $\forall a, b, c \in \mathcal{A}$, we have

$$(aD_0)(bc) = aD_0(bc) = aD_0(b)c + abD_0(c) = (aD_0)(b)c + b(aD_0)(c),$$

so aD_0 is a derivation of $\mathcal{A}$.

3) For $\forall a, b, c \in \mathcal{A}$, we have

$$\begin{aligned} [aD_0, bD_0](c) &= (aD_0)(bD_0)(c) - (bD_0)(aD_0)(c) \\ &= aD_0(b)D_0(c) - bD_0(a)D_0(c) \\ &= (aD_0(b) - bD_0(a))D_0(c). \end{aligned}$$

Then $[aD_0, bD_0] = (aD_0(b) - bD_0(a))D_0 \in \mathcal{D}_1$, and so 3) holds. □

Theorem 2.4 *Let $\mathcal{A}$ be a commutative and associative algebra satisfying Lemma 2.1, and let a be an element of $\mathcal{A}$. If D_0 satisfies Lemma 2.3 and $\circ$ satisfies*

$$b \circ c = baD_0(c), \forall b, c \in \mathcal{A}, \tag{2.4}$$

then the following statements hold:

1) $(\mathcal{A}, aD_0, \circ)$ *is a Novikov algebra.*

2) $(\mathcal{A}, aD_0, [,])$ *is an adjoining Lie algebra of* $(\mathcal{A}, aD_0, \circ)$ *and* $[\ ,\ ]$ *such that*

$$[b, c] = a(bD_0(c) - cD_0(b)), \quad \forall b, c \in \mathcal{A}. \tag{2.5}$$

Proof. 1) By Lemma 2.3, aD_0 is a derivation of the commutative algebra $\mathcal{A}$. So $(\mathcal{A}, aD_0, \circ)$ is a Novikov algebra by Remark 1.3 (2).

2) $(\mathcal{A}, aD_0, [,])$ is an adjoining Lie algebra of $(\mathcal{A}, aD_0, \circ)$ by Remark 1.3 (1). For $\forall b, c \in \mathcal{A}, \exists a \in \mathcal{A}$, we have

$$[b, c] = b \circ c - c \circ b = baD_0(c) - caD_0(b) = a(bD_0(c) - cD_0(b))$$

since $\mathcal{A}$ is commutative. Hence we obtain the desired result. □

Let b_0 be a unity of $\mathcal{A}$. If we set $a = b_0$ in Theorem 2.4, then $a_n \circ a_m = a_n b_0 D_0(a_m) = a_n(mb_m) = \frac{m}{2}(a_{m+n} + a_{n-m})$. Similarly, we obtain the following corollary:

Corollary 2.5 *Let $\mathcal{A}$ be a commutative and associative algebra satisfying Lemma 2.1. Then the following statements hold:*

$$\begin{cases} a_n \circ a_m = \frac{m}{2}(a_{n+m} + a_{n-m}) \\ b_n \circ b_m = \frac{m}{2}(a_{n+m} + a_{m-n}) \\ a_n \circ b_m = \frac{m}{2}(b_{n+m} - b_{n-m}) \\ b_n \circ a_m = \frac{m}{2}(b_{n+m} + b_{n-m}) \end{cases} \tag{2.6}$$

and

$$\begin{cases} [a_n, a_m] = \frac{1}{2}(m-n)a_{n+m} + \frac{1}{2}(m+n)a_{n-m} \\ [b_n, b_m] = \frac{1}{2}(m-n)a_{n+m} - \frac{1}{2}(m+n)a_{n-m} \\ [a_n, b_m] = \frac{1}{2}(m-n)b_{n+m} - \frac{1}{2}(n+m)b_{n-m} \\ [b_n, a_m] = \frac{1}{2}(m-n)b_{n+m} + \frac{1}{2}(m+n)b_{n-m}. \end{cases} \tag{2.7}$$

The following, let $\sinh x = \frac{e^x - e^{-x}}{2}$, $\cosh x = \frac{e^x + e^{-x}}{2}$ the field $\mathbf{F}$ be assumed $\mathbf{R}$ or $\mathbf{C}$. We will construct Novikov algebras over the linear space which is generated by $\sinh x$ and $\cosh x$.

First, let $\mathcal{T}$ be a linear space generated by $\{\sinh mx, \cosh nx | m, n \in \mathbf{N}\}$ over $\mathbf{F}$.

Lemma 2.6 *$\mathcal{T}$ satisfying the above product is a commutative associative algebra.*

Proof. Since the above product is commutative and associative, we only need that $\mathcal{T}$ is closed for the product. In fact,

$$\begin{cases} \sinh mx \sinh nx = \dfrac{1}{2}[\cosh(m+n)x - \cosh(m-n)x] \\ \\ \cosh mx \cosh nx = \dfrac{1}{2}[\cosh(m+n)x + \cosh(m-n)x] \\ \\ \sinh mx \cosh nx = \dfrac{1}{2}[\sinh(m+n)x + \sinh(m-n)x]. \end{cases} \tag{2.8}$$

So $\mathcal{T}$ is a commutative and associative algebra. □

Lemma 2.7 *Let $\mathcal{T}$ be a linear space generated by $\{\sinh mx, \cosh nx | m, n \in \mathbf{N}\}$ over $\mathbf{F}$, then $\{1, \sinh mx, \cosh nx | m, n \in \mathbf{N_0}\}$ is a basis of $\mathcal{T}$.*

Proof. For $\forall n \in \mathbf{N}_0$, suppose that there are $c_0, a_i, b_j \in \mathbf{F}, i, j \in \mathbf{N_0}$ such that

$$c_0 + a_1 \sinh x + b_1 \cosh x + \cdots + a_n \sinh nx + b_n \cosh nx = 0. \tag{2.9}$$

We take derivative for (2.9) such that its derivative order is $2k-1$ ($k \in \mathbf{N}_0$), and put $x = 0$. Then we have

$$a_1 + 2^{2k-1} a_2 + \cdots + n^{2k-1} a_n = 0.$$

Let $k = 1, 2, \cdots, n$, then we obtain the following system of n linear equations:

$$\begin{cases} a_1 + 2a_2 + \cdots + na_n = 0 \\ a_1 + 2^3 a_2 + \cdots + n^3 a_n = 0 \\ \cdots\cdots\cdots\cdots\cdots\cdots\cdots\cdots \\ a_1 + 2^{2n-1} a_2 + \cdots + n^{2n-1} a_n = 0. \end{cases} \tag{2.10}$$

If $a_1, \cdots, a_n$ are seen to be unknown, then the coefficient matrix of (2.10) is the Vandermonde matrix whose determinant is not 0, so $a_i = 0, i = 1, \cdots, n$.

We take derivative for (2.9) such that its derivative order is $2k$ ($k \in \mathbf{N}_0$), and put $x = 0$. Then we have

$$b_1 + 2^{2k} b_2 + \cdots + n^{2k} b_n = 0.$$

Let $k = 1, 2, \cdots, n$, then we obtain the following system of n linear equations:

$$\begin{cases} b_1 + 2^2 b_2 + \cdots + n^2 b_n = 0 \\ b_1 + 2^4 b_2 + \cdots + n^4 b_n = 0 \\ \cdots\cdots\cdots\cdots\cdots\cdots\cdots\cdots \\ b_1 + 2^{2n} b_2 + \cdots + n^{2n} b_n = 0. \end{cases} \tag{2.11}$$

If $b_1, \cdots, b_n$ are seen to be unknown, then the coefficient matrix of (2.11) is the Vandermonde matrix whose determinant is not 0, so $b_i = 0, i = 1, \cdots, n$. Since for $\forall i \in \mathbf{N}_0$, $a_i = 0$ and $b_i = 0$ satisfy (2.9), we have $c_0 = 0$. Hence $\{1$, $\sinh x$, $\cosh x$, $\cdots$, $\sinh nx$, $\cosh nx\}$ are linearly independent for $\forall n \in \mathbf{N}_0$, then $\{1, \sinh nx, \cosh mx | n, m \in \mathbf{N}_0\}$ are linearly independent and so they form a basis of $\mathcal{T}$ as desired. □

Theorem 2.8 *Let $\mathcal{A}_1$, $\mathcal{A}_2$ be commutative and associative algebras over $\mathbf{F}$. If φ: $\mathcal{A}_1 \longrightarrow \mathcal{A}_2$ is an isomorphism and $D_1 \in \mathrm{Der}\mathcal{A}_1$, then the following statements hold:*

1) $D_2 := \varphi D_1 \varphi^{-1} \in \mathrm{Der}\mathcal{A}_2$.

2) *φ: $(\mathcal{A}_1, D_1, \circ) \longrightarrow (\mathcal{A}_2, D_2, \circ)$ is also an isomorphism of Novikov algebras.*

Proof. 1) For $\forall a, b \in \mathcal{A}_1$, we have

$$\begin{aligned}&(\varphi D_1 \varphi^{-1})(\varphi(a)\varphi(b)) = (\varphi D_1 \varphi^{-1})(\varphi(ab))\\ &= \varphi D_1(ab) = \varphi(D_1(a)b + aD_1(b)) = \varphi(D_1(a))\varphi(b) + \varphi(a)\varphi(D_1(b))\\ &= (\varphi D_1 \varphi^{-1})(\varphi(a))\varphi(b) + \varphi(a)(\varphi D_1 \varphi^{-1})(\varphi(b)).\end{aligned}$$

So 1) holds.

2) For $\forall a, b \in \mathcal{A}_1$, we have

$$\begin{aligned}&\varphi(a \circ b) = \varphi(aD_1(b)) = \varphi(a)\varphi(D_1(b))\\ &= \varphi(a)(\varphi D_1 \varphi^{-1})(\varphi(b)) = \varphi(a)D_2(\varphi(b))\\ &= \varphi(a) \circ \varphi(b).\end{aligned}$$

So 2) holds. □

Theorem 2.9 *Let $\mathcal{A}$ be a commutative and associative algebra over $\mathbf{F}$ satisfying Lemma 2.1, D_0 be its derivation satisfying (2.2) and $\mathcal{T}$ be a commutative and associative algebra over $\mathbf{F}$ satisfying Lemmas 2.6 and 2.7. If $\varphi : \mathcal{A} \longrightarrow \mathcal{T}$ satisfies*

$$\varphi(b_m) = \cosh mx,\ m = 0, 1, 2, \cdots, \quad \varphi(a_n) = \sinh nx,\ n = 1, 2, \cdots, \tag{2.12}$$

then the following statements hold:

1) *φ is an isomorphism of commutative and associative algebras.*

2) $\varphi D_0 \varphi^{-1} = \frac{\mathrm{d}}{\mathrm{d}x}$.

3) *$\varphi : (\mathcal{A}, aD_0, \circ) \longrightarrow (\mathcal{T}, \varphi(a)\frac{\mathrm{d}}{\mathrm{d}x}, \circ)$ is an isomorphism of Novikov algebras.*

Proof. It is clear by Lemma 2.7, (2.1) and (2.8).

2) By (2.2) and (2.12), we have

$$\begin{aligned}&\varphi D_0 \varphi^{-1}(\sinh nx) = \varphi D_0(a_n)\\ &= \varphi(nb_n) = n \cosh nx\\ &= \frac{\mathrm{d} \sinh nx}{\mathrm{d}x},\\ &\varphi D_0 \varphi^{-1}(\cosh nx) = \varphi D_0(b_n)\\ &= \varphi(na_n) = n \sinh nx\\ &= \frac{\mathrm{d} \cosh nx}{\mathrm{d}x}.\end{aligned}$$

So 2) holds.

3) It is clear that $\varphi(aD_0)\varphi^{-1} = \varphi(a)d/dx$. By (2.12) and (2.2), we have

$$\begin{aligned}
&\varphi(aD_0)\varphi^{-1}(\sinh nx) = \varphi(aD_0)(a_n)\\
&= \varphi(aD_0(a_n)) = \varphi(anb_n)\\
&= \varphi(a)\varphi(nb_n) = \varphi(a)n\cosh nx\\
&= \varphi(a)d(\sinh nx)/dx.
\end{aligned}$$

Similarly, we have $\varphi(aD_0)\varphi^{-1}(\cosh nx) = \varphi(a)d(\cosh nx)/dx$. So $\varphi(aD_0)\varphi^{-1} = \varphi(a)d/dx$.

By Theorems 2.4, 2.8 and Remark 1.3 (2), we have

$$\begin{aligned}
&\varphi(b \circ c) = \varphi(baD_0(c))\\
&= \varphi(b)\varphi(aD_0(c))\\
&= \varphi(b)[\varphi(aD_0)\varphi^{-1}(\varphi(c))]\\
&= \varphi(b)\varphi(a)d/dx(\varphi(c))\\
&= \varphi(b) \circ \varphi(c), \forall b, c \in \mathcal{A}.
\end{aligned}$$

So $\varphi : (\mathcal{A}_0, aD_0, \circ) \longrightarrow (\mathcal{T}, \varphi(a)\frac{\mathrm{d}}{\mathrm{d}x}, \circ)$ is an isomorphism of Novikov algebras. $\square$

References

[1] Bai C M and Meng D J 2000 The structure of bi-symmetric algebras and their subadjacent Lie algebras *Commun. Algebra* **28** 2717-2734

[2] Bai C M and Meng D J 2001 The classification of Novikov algebras in low dimensions *J. Phys. A: Math. Gen.* **34** 1581-1594

[3] Bai C M and Meng D J 2001 The realizations of non-transitive Novikov algebrss *J. Phys. A: Math. Gen.* **34** 6435-6442

[4] Chen L Y, Niu Y J and Meng D J 2008 Two kinds of Novikov algebras and their realizations *J. Pure Appl. Algebra* **212** 902-909

[5] Gel'fand I M and Diki L A 1975 Asymptotic behavior of the resolvent of Sturm-Liou-ville equations and the Lie algebra of the Korteweg-de Vries equations *Russian Math. Surveys* **30** 77-113

[6] Gel'fand I M and Diki L A 1976 A Lie algebra structure in a formal variational cal-culation *Functional Anal. Appl.* **10** 16-22

[7] Gel'fand I M and Dorfman I Y 1979 Hamiltonian operators and algebraic structures related to them *Functional Anal. Appl.* **13** 248-262

[8] Kang Y F and Chen Z Q 2009 Novikov superalgebras in low dimensions *J. Nonlinear math.Phy.* **16(3)** 251-257

[9] Osborn J M 1992 simple Novikov algebras with an idempotent *Commun. Algebra* **20** 2729-2753

[10] Osborn J M 1994 Infinite dimentional Novikov algebras of characteristic 0 *J. Algebra* **16** 146-167

[11] Xu X P 1995 Hamiltonian operators and associative algebras whih a derivation *Lett. Math. Phys.* **33** 1-6

[12] Xu X P 1996 On simple Novikov algebra and their irreducible modules *J. Algebra* **185** 905-934

[13] Xu X P 1997 Novikov-Poisson algebras *J. Algebra* **190** 253-279

[14] Xu X P 2000 Variational calculus of supervarables and related algebraic structures *J. Algebra* **223** 396-437

[15] Zel'manov E I 1987 On a class of local translation invariant Lie algebras *Soviet Math. Dokl.* **35(1)** 216-218

[16] Zhu F H and Chen Z Q 2010 Novikov superalgebras with $A_0 = A_1A_1$ *Czechoslovak Mathematical Journal* **60(135)** 903-907